

Gauge Theories from Orientifolds and Large N Limit

Zurab Kakushadze*

Lyman Laboratory of Physics, Harvard University, Cambridge, MA 02138

and

Department of Physics, Northeastern University, Boston, MA 02115

(March 25, 1998)

Abstract

Extending the recent work in [1], we consider string perturbative expansion in the presence of D-branes and orientifold planes imbedded in orbifolded space-time. In the $\alpha' \rightarrow 0$ limit the weak coupling string perturbative expansion maps to 't Hooft's large N expansion. We focus on four dimensional $\mathcal{N} = 1, 2, 4$ supersymmetric theories, and also discuss possible extensions to $\mathcal{N} = 0$ cases. Utilizing the string theory perturbation techniques we show that computation of any M -point correlation function in these theories reduces to the corresponding computation in the parent $\mathcal{N} = 4$ theory. In particular, we discuss theories (which are rather constrained) with vanishing β -functions to all orders in perturbation theory in the large N limit. We also point out that in theories with non-vanishing β -functions the gauge coupling running is suppressed in the large N limit. Introduction of orientifold planes allows to construct certain gauge theories with SO , Sp and SU gauge groups and various matter (only unitary gauge groups with bi-fundamental/adjoint matter arise in theories without orientifold planes).

Typeset using REVTeX

*E-mail: zurab@string.harvard.edu

I. INTRODUCTION

‘t Hooft’s large N expansion [2] is an attractive possibility for understanding gauge theories. In this limit the gauge theory diagrams look like Riemann surfaces with boundaries and handles. It is therefore natural to attempt to map the large N expansion of gauge theories to some kind of string world-sheet expansion.

The first concrete example of such a map was given by Witten [3] for the case of three dimensional Chern-Simons gauge theory where the boundaries of the string world-sheet are “topological” D-branes.

Recently the map between the large N expansion and string expansion has been made precise [1] in the context of Type II string theory. The idea of [1] is to consider Type IIB string theory with a large number N of D3-branes and take a limit where $\alpha' \rightarrow 0$ while keeping $\lambda = N\lambda_s$ fixed, where λ_s is the Type IIB string coupling. In this setup we have four dimensional gauge theories with unitary gauge groups. A world-sheet with b boundaries and g handles is weighted with

$$(N\lambda_s)^b \lambda_s^{2g-2} = \lambda^{2g-2+b} N^{-2g+2} . \quad (1)$$

Upon identification $\lambda_s = g_{YM}^2$, this precisely maps to ‘t Hooft’s large N expansion. This expansion is valid in the limit where $N \rightarrow \infty$ and the effective coupling λ is fixed at a weak coupling value.

In [1] the above idea was applied to prove that four dimensional gauge theories (including the cases with no supersymmetry) considered in [4,5]¹ are conformal to all orders in perturbation theory in the large N limit. In particular, the corresponding gauge theories were obtained from Type IIB string theory with D3-branes imbedded in orbifolded space-time. The ultraviolet finiteness of string theory (that is, one-loop tadpole cancellation conditions) was shown to imply that the resulting (non-Abelian) gauge theories were conformal in the large N limit (in all loop orders). Moreover, in [1] it was also proven that computation of any correlation function in these theories in the large N limit reduces to the corresponding computation in the parent $\mathcal{N} = 4$ supersymmetric gauge theory.

The all-order proofs in [1] were possible due to the fact that the power of string perturbation techniques was utilized. In particular, string theory perturbation expansion is a very efficient way of summing up various field theory diagrams. Thus, often a large number of field theory Feynman diagrams in a given order of perturbation theory corresponds to a single string theory diagram with certain topology. This has been successfully exploited to compute tree and loop level scattering processes in gauge theories [8]. (For a recent discussion, see, *e.g.*, [9].) The all-order proofs in [1] crucially depended on the fact that the string world-sheet expansion was self-consistent. In particular, the arguments in [1] would not go through if the tadpoles were not cancelled. The tadpole cancellation conditions, however, ultimately produced theories which were (super)conformal in the large N limit. In particular, the one-loop β -functions (for non-Abelian gauge groups) in all of those theories were zero (even at finite N).

¹For other related works, see, *e.g.*, [6,7].

It is natural to ask whether there exists a consistent string theory setup similar to that of [1] which can produce: (i) finite gauge theories with gauge groups other than unitary (in particular, SO and Sp), and with matter other than bi-fundamentals; (ii) non-finite gauge theories such that string theory perturbation techniques can be applied to learn about the gauge theories in the large N limit. In this paper we consider precisely such a setup which is a generalization of the work in [1]. Here we will consider Type IIB string theory with D3-branes as well as orientifold planes. (In certain cases string consistency will also require presence of D7-branes.) That is, we are going to study gauge theories that arise from Type IIB orientifolds. In this framework we can expect appearance of SO and Sp gauge groups as well as matter other than bi-fundamentals (in the product gauge groups). Moreover, as we will see later, non-finite gauge theories can also be obtained within a consistent perturbative string theory framework. However, in all such theories obtained via orientifolds the running of the gauge couplings is suppressed in the large N limit. (In this sense these theories are “finite” in the large N limit).

Introduction of orientifold planes changes the possible topologies of the world-sheet. Now we can have a world-sheet with b boundaries (corresponding to D-branes), c cross-caps (corresponding to orientifold planes), and g handles. Such a world-sheet is weighted with

$$(N\lambda_s)^b \lambda_s^c \lambda_s^{2g-2} = \lambda^{2g-2+b+c} N^{-c-2g+2} . \quad (2)$$

Note that addition of a cross-cap results in a diagram suppressed by an additional power of N , so that in the large N limit we can hope for simplifications (or, rather, we can hope to avoid complications with unoriented world-sheets, at least in some cases). In fact, we will show that for string vacua which are perturbatively consistent (that is, the tadpoles cancel) calculations of correlation functions in $\mathcal{N} < 4$ gauge theories reduce to the corresponding calculations in the parent $\mathcal{N} = 4$ *oriented* theory. This holds not only for finite (in the large N limit) gauge theories but also for the gauge theories which are not conformal. We will prove that certain $\mathcal{N} = 1$ gauge theories obtained this way are superconformal to all orders in perturbation theory in the large N limit. We will also discuss possible extensions to non-supersymmetric cases.

It is very satisfying to observe that in the large N limit using the power of string theory perturbation techniques we can reduce very non-trivial calculations in gauge theories with lower supersymmetries to calculations in $\mathcal{N} = 4$ gauge theories. In particular, this applies to multi-point correlators in gauge theory.

Here we note that the correspondence between ‘t Hooft’s large N expansion and string world-sheet expansion is expected to hold only in the regime where the effective coupling λ is small. If λ is large one expects an effective supergravity description to take over. This domain has been recently studied in several papers [10–15] as well as in various related works [16]. The supergravity picture has, in particular, led to the conjectures in [4] as well as in [5] about finiteness of certain gauge theories. However, proofs of those conjectures (in the large N limit) presented in [1] were given in the *weakly* coupled region. Also, $1/N$ corrections can only be reliably computed in this region but not in the strong coupling regime where *a priori* there is no world-sheet expansion nor ‘t Hooft’s expansion is valid.

The remainder of this paper is organized as follows. In section II we review the arguments of [1] for the cases without the orientifold planes. In section III we generalize these arguments to the orientifold cases. In particular, we prove various vanishing theorems in the large N

limit, and give the relation between various correlators in $\mathcal{N} < 4$ gauge theories to those in the parent $\mathcal{N} = 4$ theory. In section IV we construct some explicit $\mathcal{N} = 2$ examples. All of these examples turn out to be superconformal. In section V we construct an example of $\mathcal{N} = 1$ gauge theory with $Sp(N)$ gauge group. This gauge theory is superconformal in the large N limit to all orders in perturbation theory. We also discuss some examples of non-finite $\mathcal{N} = 1$ gauge theories.

II. LARGE N LIMIT AND FINITENESS

In this section we review the discussion in [1] for the cases without orientifold planes. We will generalize these arguments to the orientifold cases in section III.

A. Setup

Consider Type IIB string theory with N parallel D3-branes where the space transverse to the D-branes is $\mathcal{M} = \mathbf{R}^6/\Gamma$. The orbifold group $\Gamma = \{g_a \mid a = 1, \dots, |\Gamma|\} (g_1 = 1)$ must be a finite discrete subgroup of $Spin(6)$. If $\Gamma \subset SU(3)$ ($SU(2)$), we have $\mathcal{N} = 1$ ($\mathcal{N} = 2$) unbroken supersymmetry, and $\mathcal{N} = 0$, otherwise.

Let us confine our attention to the cases where type IIB on $\mathcal{M}$ is a modular invariant theory². The action of the orbifold on the coordinates X_i ($i = 1, \dots, 6$) on $\mathcal{M}$ can be described in terms of $SO(6)$ matrices: $g_a : X_i \rightarrow \sum_j (g_a)_{ij} X_j$. We need to specify the action of the orbifold group on the Chan-Paton charges carried by the D3-branes. It is described by $N \times N$ matrices γ_a that form a representation of Γ . Note that γ_1 is an identity matrix and $\text{Tr}(\gamma_1) = N$.

The D-brane sector of the theory is described by an *oriented* open string theory. In particular, the world-sheet expansion corresponds to summing over oriented Riemann surfaces with arbitrary genus g and arbitrary number of boundaries b , where the boundaries of the world-sheet correspond to the D3-branes.

For example, consider one-loop vacuum amplitude ($g = 0$, $b = 2$). The corresponding graph is an annulus whose boundaries lie on D3-branes. The one-loop partition function in the light-cone gauge is given by

$$\mathcal{Z} = \frac{1}{2|\Gamma|} \sum_a \text{Tr} \left(g_a (1 + (-1)^F) e^{-2\pi t L_0} \right), \quad (3)$$

where F is the fermion number operator, t is the real modular parameter of the annulus, and the trace includes sum over the Chan-Paton factors.

The orbifold group Γ acts on both ends of the open strings. The action of $g_a \in \Gamma$ on Chan-Paton charges is given by $\gamma_a \otimes \gamma_a$. Therefore, the individual terms in the sum in (3) have the following form:

²This is always the case if $\Gamma \subset SU(3)$. For the non-supersymmetric cases this is also true provided that $\nexists \mathbf{Z}_2 \subset \Gamma$. If $\exists \mathbf{Z}_2 \subset \Gamma$, then modular invariance requires that the set of points in $\mathbf{R}^6$ fixed under the $\mathbf{Z}_2$ twist has real dimension 2.

$$(\text{Tr}(\gamma_a))^2 \mathcal{Z}_a , \quad (4)$$

where $\mathcal{Z}_a$ are characters corresponding to the world-sheet degrees of freedom. The “un-twisted” character $\mathcal{Z}_1$ is the same as in the $\mathcal{N} = 4$ theory for which $\Gamma = \{1\}$. The information about the fact that the orbifold theory has reduced supersymmetry is encoded in the “twisted” characters $\mathcal{Z}_a$, $a \neq 1$.

In [1] it was shown that the one-loop massless (and, in non-supersymmetric cases, tachyonic) tadpole cancellation conditions require that

$$\text{Tr}(\gamma_a) = 0 \quad \forall a \neq 1 . \quad (5)$$

It was also shown that this condition implies that the Chan-Paton matrices γ_a form an n -fold copy of the *regular* representation of Γ . The regular representation decomposes into a direct sum of all irreducible representations $\mathbf{r}_i$ of Γ with degeneracy factors $n_i = |\mathbf{r}_i|$. The gauge group is ($N_i \equiv nn_i$)

$$G = \otimes_i U(N_i) . \quad (6)$$

The matter consists of Weyl fermions (and scalars) transforming in bi-fundamentals $(\mathbf{N}_i, \overline{\mathbf{N}}_j)$ according to the decomposition of the tensor product of $\mathbf{4}$ ($\mathbf{6}$) of $Spin(6)$ with the corresponding representation (see [5] for details).

B. Large N Limit

The gauge group in the theories we are considering here is $G = \otimes_i U(N_i) (\subset U(N))$. In the following we will ignore the $U(1)$ factors (for which the gauge couplings do run for $\mathcal{N} < 4$ as there are matter fields charged under them) and consider $G = \otimes_i SU(N_i)$. In this subsection we review the arguments of [1] which show that in the large N limit this non-Abelian gauge theory is conformal³.

There are two classes of diagrams we need to consider: (i) diagrams without handles; (ii) diagrams with handles. The latter correspond to closed string loops and are subleading in the large N limit. The diagrams without handles can be divided into two classes: (i) planar diagrams where all the external lines are attached to the same boundary; (ii) non-planar diagrams where the external lines are attached to at least two different boundaries. The latter are subleading in the large N limit.

In the case of planar diagrams we have b boundaries with all M external lines attached to the same boundary (which without loss of generality can be chosen to be the outer boundary) as depicted in Fig.1. We need to sum over all possible twisted boundary conditions for the boundaries. The boundary conditions must satisfy the requirement that

$$\gamma_{a_1} = \prod_{s=2}^b \gamma_{a_s} , \quad (7)$$

³Including the $U(1)$ factors does not alter the conclusions as their effect is subleading in the $1/N$ expansion [1].

where γ_{a_1} corresponds to the outer boundary (to which we have attached the external lines), and γ_{a_s} ($s = 2, \dots, b$) correspond to the inner boundaries (with no external lines). Here we have chosen the convention (consistent with the corresponding choice made for the annulus amplitude in (3)) that the outer and inner boundaries have opposite orientations. Then the above condition is simply the statement that only the states invariant under the action of the orbifold group contribute into the amplitude.

If all the twisted boundary conditions are trivial (*i.e.*, $a_s = 1$ for all $s = 1, \dots, b$) then the corresponding amplitude is the same as in the $\mathcal{N} = 4$ case (modulo factors of $1/\sqrt{|\Gamma|}$ coming from the difference in normalizations of the corresponding D-brane boundary states in the cases with $\mathcal{N} = 4$ (where $|\Gamma| = 1$) and $\mathcal{N} < 4$ (where $|\Gamma| \neq 1$)). Therefore, such amplitudes do not contribute to the gauge coupling running (for which we would have $M = 2$ gauge bosons attached to the outer boundary) since the latter is not renormalized in $\mathcal{N} = 4$ gauge theories due to supersymmetry.

Let us now consider contributions with non-trivial twisted boundary conditions. Let λ_r , $r = 1 \dots M$, be the Chan-Paton matrices corresponding to the external lines. Then the planar diagram with b boundaries has the following Chan-Paton group-theoretic dependence:

$$\sum \text{Tr}(\gamma_{a_1} \lambda_1 \dots \lambda_M) \prod_{s=2}^b \text{Tr}(\gamma_{a_s}) , \quad (8)$$

where the sum involves all possible distributions of γ_{a_s} twists (that satisfy the condition (7)) as well as permutations of λ_r factors (note that the λ 's here are the states which are kept after the orbifold projection, and so they commute with the action of γ 's). The important point here is that unless all twists γ_{a_s} are trivial for $s = 2, \dots, b$, the above diagram vanishes by the virtue of (5). But then from (7) it follows that γ_{a_1} must be trivial as well. This implies that the only planar diagrams that contribute are those with trivial boundary conditions which (up to numerical factors) are the same as in the parent $\mathcal{N} = 4$ gauge theory. This establishes that computation of any M -point correlation function in the large N limit reduces to the corresponding $\mathcal{N} = 4$ calculation, and that these gauge theories are (super)conformal in this limit⁴.

Here we should mention that the models of [1] are perturbatively consistent string theories at all energy scales. In particular, the Abelian factors (that run in the low energy effective theory and decouple in the infrared) are not problematic from the string theory viewpoint (although in the field theory context they would have Landau poles in the ultraviolet).

III. GENERALIZATION TO ORIENTIFOLDS

In this section we generalize the approach of [1] to theories with orientifold planes. Here we will mostly concentrate on supersymmetric cases, and briefly discuss possible generalizations to non-supersymmetric cases at the end of this section.

⁴In [1] it was also shown that a large class of non-planar diagrams without handles also vanish. We refer the reader to [1] for details.

A. Setup

Consider Type IIB string theory on $\mathcal{M} = \mathbf{C}^3/\Gamma$ where $\Gamma \subset SU(3)$ so that the resulting theory has some number of unbroken supersymmetries. Consider the ΩJ orientifold of this theory, where Ω is the world-sheet parity reversal, and J is a $\mathbf{Z}_2$ element ($J^2 = 1$) acting on the complex coordinates z_i ($i = 1, 2, 3$) on $\mathbf{C}^3$ such that the set of points in $\mathbf{C}^3$ fixed under the action of J has real dimension $\Delta = 0$ or 4.

If $\Delta = 0$ then we have an orientifold 3-plane. If Γ has a $\mathbf{Z}_2$ subgroup, then we also have an orientifold 7-plane. If $\Delta = 4$ then we have an orientifold 7-plane. We may also have an orientifold 3-plane depending on whether Γ has an appropriate $\mathbf{Z}_2$ subgroup. Regardless of whether we have an orientifold 3-plane, we can *a priori* introduce an arbitrary number of D3-branes (as the corresponding tadpoles vanish due to the fact that the space transverse to the D3-branes is non-compact). On the other hand, if we have an orientifold 7-plane we must introduce 8 of the corresponding D7-branes to cancel the R-R charge appropriately. (The number 8 of D7-branes is required by the corresponding tadpole cancellation conditions.)

We need to specify the action of Γ on the Chan-Paton factors corresponding to the D3- and/or D7-branes. Just as in the previous section, these are given by Chan-Paton matrices which we collectively refer to as γ_a^μ , where the superscript μ refers to the corresponding D3- or D7-branes. Note that $\text{Tr}(\gamma_1^\mu) = n^\mu$ where n^μ is the number of D-branes labelled by μ .

At one-loop level there are three different sources for massless tadpoles: the Klein bottle, annulus, and Möbius strip amplitudes depicted in Fig.2. The Klein bottle amplitude corresponds to the contribution of unoriented closed strings into one-loop vacuum diagram. It can be alternatively viewed as a tree-level closed string amplitude where the closed strings propagate between two cross-caps. The latter are (coherent Type IIB) states that describe the familiar orientifold planes. The annulus amplitude corresponds to the contribution of open strings stretched between two D-branes into one-loop vacuum amplitude. It can also be viewed as a tree-channel closed string amplitude where the closed strings propagate between two D-branes. Finally, the Möbius strip amplitude corresponds to the contribution of unoriented open strings into one-loop vacuum diagram. It can be viewed as a tree-channel closed string amplitude where the closed strings propagate between a D-brane and an orientifold plane.

Note that there are no Chan-Paton matrices associated with the Klein bottle amplitude since it corresponds to closed strings propagating between two cross-caps which do not carry Chan-Paton charges. The Möbius strip has only one boundary. This implies that the individual terms (corresponding to twists $g_a \in \Gamma$) in the Möbius strip amplitude are proportional to $\text{Tr}(\gamma_a^\mu)$. The annulus amplitude is the same (up to an overall factor of 1/2 due to the orientation reversal projection) as in the oriented case discussed in the previous section. Thus, the individual terms (corresponding to twists $g_a \in \Gamma$) in the annulus amplitude are proportional to $\text{Tr}(\gamma_a^\mu)\text{Tr}(\gamma_a^\nu)$. Thus, the tadpoles can be written as

$$\sum_a \left(K_a + \sum_\mu M_a^\mu \text{Tr}(\gamma_a^\mu) + \sum_{\mu,\nu} A_a^{\mu\nu} \text{Tr}(\gamma_a^\mu) \text{Tr}(\gamma_a^\nu) \right). \quad (9)$$

Here terms with K_a , M_a^μ and $A_a^{\mu\nu}$ correspond to the contributions of the Klein bottle, Möbius strip and annulus amplitudes, respectively. In fact, the factorization property of string theory implies that the Klein bottle amplitude should factorize into two cross-caps

connected via a long thin tube. The Möbius strip amplitude should factorize into a cross-cap and a disc connected via a long thin tube. Similarly, the annulus amplitude should factorize into two discs connected via a long thin tube. These factorizations are depicted in Fig.3. The implication of this for the tadpoles is that they too factorize into a sum of perfect squares

$$\sum_a \left(B_a + \sum_\mu C_a^\mu \text{Tr}(\gamma_a^\mu) \right)^2, \quad (10)$$

where $B_a^2 = K_a$, $2B_a C_a^\mu = M_a^\mu$ and $C_a^\mu C_a^\nu = A_a^{\mu\nu}$. Thus, the tadpole cancellation conditions read:

$$B_a + \sum_\mu C_a^\mu \text{Tr}(\gamma_a^\mu) = 0. \quad (11)$$

Note that

$$\text{Tr}(\gamma_a^\mu) = 0 \quad \forall a \neq 1 \text{ only if } K_a = 0 \quad \forall a \neq 1. \quad (12)$$

In the next subsection we will see that if this condition is satisfied then the corresponding (non-Abelian) gauge theories are superconformal in the large N limit. On the other hand, if not all K_a with $a \neq 1$ are zero, then some of the Chan-Paton matrices γ_a^μ with $a \neq 1$ must have non-zero traces. This will generically lead to theories with non-vanishing one-loop β -functions.

B. Large N Limit

In this subsection we extend the arguments reviewed in section II to the cases with orientifold planes. In particular, we will study the large N behavior of M -point correlators of fields charged under the gauge group that arises from the D3-branes. In the following we will ignore the $U(1)$ factors (if any) in the D3-brane gauge group.

There are two classes of diagrams we need to consider: (i) diagrams without handles and cross-caps; (ii) diagrams with handles and/or cross-caps. The latter are subleading in the large N limit. The diagrams without handles and cross-caps can be divided into two classes: (i) planar diagrams where all the external lines are attached to the same boundary; (ii) non-planar diagrams where the external lines are attached to at least two different boundaries. The latter are subleading in the large N limit.

In the case of planar diagrams we have b boundaries corresponding to D3- and/or D7-branes. We will attach M external lines to the outer boundary as depicted in Fig.1. We need to sum over all possible twisted boundary conditions for the boundaries. The boundary conditions must satisfy the requirement that

$$\gamma_{a_1}^{\mu_1} = \prod_{s=2}^b \gamma_{a_s}^{\mu_s}, \quad (13)$$

where $\gamma_{a_1}^{\mu_1}$ corresponds to the outer boundary, and $\gamma_{a_s}^{\mu_s}$ ($s = 2, \dots, b$) correspond to the inner boundaries.

Let us first consider the $\mathcal{N} = 4$ theories for which the orbifold group Γ is trivial. (Note that in this case we can only have D3-branes as introduction of D7-branes would break some number of supersymmetries.) The computation of any correlation function in the orientifold theory (with $SO(N)$ or $Sp(N)$ gauge group) is reduced to the corresponding computation in the oriented $\mathcal{N} = 4$ theory (with $U(N)$ gauge group) up to factors of $1/\sqrt{2}$ (coming from the difference in normalizations of the corresponding D-brane boundary states in the oriented and unoriented cases). Such a simplification is due to the fact that the unoriented world-sheets with cross-caps give contributions suppressed by extra powers of $1/N$.

Next, consider cases where Γ is non-trivial (and hence supersymmetry is reduced). If all the twisted boundary conditions are trivial (*i.e.*, $a_s = 1$ for all $s = 1, \dots, b$) then the corresponding amplitude is the same as in the $\mathcal{N} = 4$ case (modulo factors of $1/\sqrt{|\Gamma|}$). Therefore, such amplitudes do not contribute to the gauge coupling running (for which we would have $M = 2$ gauge bosons attached to the outer boundary) since the latter is not renormalized in $\mathcal{N} = 4$ gauge theories due to supersymmetry.

Let us now consider contributions with non-trivial twisted boundary conditions. Let λ_r , $r = 1 \dots M$, be the Chan-Paton matrices corresponding to the external lines. Then the planar diagram with b boundaries has the following Chan-Paton group-theoretic dependence:

$$\sum \text{Tr} \left(\gamma_{a_1}^{\mu_1} \lambda_1 \dots \lambda_M \right) \prod_{s=2}^b \text{Tr}(\gamma_{a_s}^{\mu_s}) , \quad (14)$$

where the sum involves all possible distributions of γ_{a_s} twists (that satisfy the condition (7)) as well as permutations of λ_r factors. If the condition (12) is satisfied, *i.e.*, if all the twisted Chan-Paton matrices are traceless, then the situation is analogous to that in the oriented cases. That is, the only planar diagrams that contribute are those with trivial boundary conditions. Such diagrams with all the boundaries corresponding to D3-branes (up to numerical factors) are the same as in the parent $\mathcal{N} = 4$ gauge theory. The diagrams with trivial boundary conditions but with some boundaries corresponding to D7-branes are subleading in the large N limit as the numbers of D7-branes (that is, the traces $\text{Tr}(\gamma_1^\mu)$ corresponding to D7-branes) are of order one. This establishes that computation of any M -point correlation function in the large N limit reduces to the corresponding $\mathcal{N} = 4$ calculation in *oriented* theory, and that these gauge theories are superconformal in this limit⁵.

Now consider the cases where some of the twisted Chan-Paton matrices are *not* traceless. Then there are going to be corrections to the M -point correlators coming from planar diagrams with non-trivial twisted boundary conditions. These diagrams are subleading in the large N limit as the corresponding traces are always of order one. This follows from the tadpole cancellation conditions (10) where the coefficients B_a and C_a^μ are of order one, so for $a \neq 1$ we have $\text{Tr}(\gamma_a^\mu) \sim 1$. This implies that even for non-finite theories computation of the correlation functions reduces to the corresponding computation in the parent $\mathcal{N} = 4$ oriented theory.

Here we should point out that “non-finiteness” of such theories is a subleading effect in the large N limit. This is because the β -function coefficients grow as

⁵Just as in [1], it is also straightforward to show that a large class of non-planar diagrams without handles and cross-caps also vanish.

$$b_s = O(N^s) , \quad s = 0, 1, \dots , \quad (15)$$

instead of $b_s = O(N^{s+1})$ (as in, say, pure $SU(N)$ gauge theory). This can be seen by considering planar diagrams with $M = 2$ external lines corresponding to gauge bosons. Note that in string theory running of the gauge couplings in the low energy effective field theory is due to infrared divergences corresponding to massless modes [17]. The diagrams with all the boundaries corresponding to D3-branes and with all the boundary conditions corresponding to the identity element of Γ are the same (up to overall numerical factors) as in the parent $\mathcal{N} = 4$ theory. Such diagrams, therefore, do not contain infrared divergencies, and thus do not contribute the gauge coupling running. (That is, their contributions to the β -function coefficients b_s vanish.) Therefore, the only diagrams that can contribute to the β -function coefficients b_s are those with some boundaries corresponding to D7-branes and/or having twisted boundary conditions with $\text{Tr}(\gamma_a^\mu) \neq 0$. These are, however, suppressed at least by one power of N since the numbers of D7-branes are of order one, and also such $\text{Tr}(\gamma_a^\mu) \sim 1$. This establishes (15).

Note that the estimates for the β -function coefficients in (15) for $b_{s>0}$ are non-trivial from the field theory point of view as they imply infinitely many cancellations between couplings (such as Yukawas) in the gauge theory. On the other hand, within string perturbation expansion these statements become obvious once we carefully consider twisted boundary conditions and tadpole cancellation.

C. Comments

Here we would like to comment on some issues concerning the discussion in the previous subsections.

First, note that the entire argument in the previous subsection crucially depends on the assumption that there is a well defined world-sheet description of the orientifold theories at hand. Naively, it might seem that orientifolds of Type IIB on $\mathbf{C}^3/\Gamma$ should have such world-sheet descriptions for any Γ which is a subgroup of $Spin(6)$. This is, however, not the case [18]⁶. In particular, there are certain cases where perturbative description is inadequate as there are additional states present in the corresponding orientifolds such that they are non-perturbative from the orientifold viewpoint [18]. In section V we will give an explicit example of this. For further details we refer the reader to [18].

The second comment is on possible extensions of the above results to $\mathcal{N} = 0$ theories. Note that in the cases without orientifold planes [1] the tadpole cancellation conditions for both $\mathcal{N} = 0$ and supersymmetric theories are given by (5) for which there always exists a solution for any choice of Γ . The situation in the cases with orientifold planes might not be so simple: the tadpoles are of the form (9), and the coefficients $K_a, M_a^\mu, A_a^{\mu\nu}$ may be different for the massless and tachyonic tadpoles. (The latter are absent in the supersymmetric cases but are generically present for $\mathcal{N} = 0$.) It is *a priori* unclear whether the massless and tachyonic tadpole cancellation conditions are compatible. However, if they have a solution in a given model, then the statements about the correlation functions in subsection B persist

⁶Some examples of this were also given in [19].

for such $\mathcal{N} = 0$ models in the large N limit. It would therefore be interesting to construct $\mathcal{N} = 0$ models in which all the tadpoles cancel.

IV. $\mathcal{N} = 2$ SUPERCONFORMAL GAUGE THEORIES

In this section we construct four dimensional $\mathcal{N} = 2$ superconformal gauge theories from Type IIB orientifolds. We start from Type IIB string theory on $\mathcal{M} = \mathbf{C}^3/\Gamma$, where $\Gamma = \{g^k | k = 0, \dots, M-1\} \approx \mathbf{Z}_M$ is the orbifold group whose action on the complex coordinates z_i ($i = 1, 2, 3$) on $\mathbf{C}^3$ is given by $gz_1 = z_1$, $gz_2 = \omega z_2$, $gz_3 = \omega^{-1} z_3$ ($\omega = \exp(2\pi i/M)$). Next, we consider an orientifold of this theory where the orientifold action is given by ΩJ . Here Ω is the world-sheet parity reversal, and J acts on z_i as $Jz_i = -z_i$. The orientifold group is given by $\mathcal{O} = \{g^k, \Omega J g^k | k = 0, \dots, M-1\}$.

A. Orbifolds of Even Order

Let us first consider the cases where M is even. Then the ΩJ orientifold of Type IIB on $\mathcal{M}$ is equivalent to the $\Omega \tilde{J}$ orientifold of Type IIB on $\mathcal{M}$, where $\tilde{J}$ acts on z_i as $\tilde{J}z_1 = -z_1$, $\tilde{J}z_2 = z_2$, $\tilde{J}z_3 = z_3$. (This is due to the presence of the $\mathbf{Z}_2$ element R in Γ where $Rz_1 = z_1$, $Rz_2 = -z_2$, $Rz_3 = -z_3$. Note that $\tilde{J} = JR$.) These orientifolds are similar to those considered in [20–22]. In particular, calculation of massless tadpoles closely parallels that of [22].

The orientifolds (with $M \in 2\mathbf{N}$) we are considering here contain both D3- and D7-branes. The action of the orbifold group on the Chan-Paton charges carried by the D3- and D7-branes is described by matrices $\gamma_{k,3}$ and $\gamma_{k,7}$ that form a (projective) representation⁷ of Γ . Note that $\gamma_{0,3}$ and $\gamma_{0,7}$ are identity matrices, and $\text{Tr}(\gamma_{0,3}) = n_3$ and $\text{Tr}(\gamma_{0,7}) = n_7$, where n_3 and n_7 are the numbers of D3- and D7-branes, respectively.

The massless tadpoles can be computed following [22]. There are two types of massless tadpoles we need to consider. The first type of tadpoles correspond to untwisted closed string exchange in the tree-channel. Cancellation of untwisted tadpoles implies that there are $n_7 = 8$ D7-branes present (for even M). On the other hand, the number of D3-branes is unconstrained by the untwisted tadpole cancellation conditions. This is due to the fact that the space transverse to the D3-branes is non-compact in the theories we are considering here.

Next, consider the tadpoles corresponding to twisted closed string exchange in the tree-channel. For even M the twisted massless tadpole cancellation conditions read:

$$\text{Tr}(\gamma_{2k-1,7}) - 4 \sin^2 \left(\frac{(2k-1)\pi}{M} \right) \text{Tr}(\gamma_{2k-1,3}) = 0 , \quad (16)$$

$$\text{Tr}(\gamma_{2k,7}) - 4 \sin^2 \left(\frac{2\pi k}{M} \right) \text{Tr}(\gamma_{2k,3}) - 8 \cos \left(\frac{2\pi k}{M} \right) = 0 . \quad (17)$$

⁷A *priori* they can form a representation of the double cover of Γ .

Note that the difference between these tadpole cancellation conditions and those of [22] is that in the second line we have $8 \cos(2\pi k/M)$ instead of $32 \cos(2\pi k/M)$. This is due to the fact that we are considering a system of D3- and D7-branes instead of a system of D5- and D9-branes⁸.

It is not difficult to show that the above tadpole cancellation conditions have solutions⁹ only for $M = 2, 4, 6$. The corresponding Chan-Paton matrices (up to equivalent representations) are given by:

- $M = 2$ ($N = n_3/2$):

$$\gamma_{1,3} = \text{diag}(i \text{ (} N \text{ times)}, -i \text{ (} N \text{ times)}) , \quad (18)$$

$$\gamma_{1,7} = \text{diag}(i \text{ (} 4 \text{ times)}, -i \text{ (} 4 \text{ times)}) . \quad (19)$$

- $M = 4$ ($N = n_3/4$):

$$\gamma_{1,3} = \text{diag}(\exp(\pi i/4) \text{ (} N \text{ times)}, \exp(-\pi i/4) \text{ (} N \text{ times)}, \\ \exp(3\pi i/4) \text{ (} N \text{ times)}, \exp(-3\pi i/4) \text{ (} N \text{ times)}) , \quad (20)$$

$$\gamma_{1,7} = \text{diag}(\exp(\pi i/4) \text{ (} 2 \text{ times)}, \exp(-\pi i/4) \text{ (} 2 \text{ times)}, \\ \exp(3\pi i/4) \text{ (} 2 \text{ times)}, \exp(-3\pi i/4) \text{ (} 2 \text{ times)}) . \quad (21)$$

- $M = 6$ ($N = (n_3 - 2)/6$):

$$\gamma_{1,3} = \text{diag}(i \exp(2\pi i/3) \text{ (} N \text{ times)}, -i \exp(2\pi i/3) \text{ (} N \text{ times)}, \\ i \exp(-2\pi i/3) \text{ (} N \text{ times)}, -i \exp(-2\pi i/3) \text{ (} N \text{ times)}), \\ i \text{ ((} N + 1 \text{) times)}, -i \text{ ((} N + 1 \text{) times)}) , \quad (22)$$

$$\gamma_{1,7} = \text{diag}(i \exp(2\pi i/3), -i \exp(2\pi i/3), i \exp(-2\pi i/3), -i \exp(-2\pi i/3), \\ i \text{ (} 2 \text{ times)}, -i \text{ (} 2 \text{ times)}) . \quad (23)$$

The massless spectra (including twisted closed string sectors) of these models are given in Table I.

⁸The former can be obtained from the latter by compactifying on T^2 , T-dualizing, and taking the dimensions of the dual torus $\tilde{T}^2$ to infinity. The orientifold p -planes ($p = 5, 9$) split into 4 orientifold $(p - 2)$ -planes upon T-dualizing. These are located at four fixed points of $\tilde{T}^2/\mathbf{Z}_2$. However, after taking the dimensions of $\tilde{T}^2$ to infinity (*i.e.*, when considering $\mathbf{C}/\mathbf{Z}_2$ instead of $\tilde{T}^2/\mathbf{Z}_2$), only one fixed point (at the origin) remains. This results in reduction of all the tadpoles by a factor of 4. In particular, the number of D7-branes is 8 as opposed to 32 in the case of D9-branes.

⁹Here we must take into account that the Chan-Paton matrices $\gamma_{k,3}$ and $\gamma_{k,7}$ must form a (projective) representation of the orbifold group Γ (or its double cover). Moreover, the orientifold projection Ω for the D-branes must be such that before taking into account the orbifold projections the gauge groups coming from the D3- and D7-branes must be Sp and SO , respectively [20,21].

Note that the one-loop β -functions for the non-Abelian subgroups of the 33 open string sector gauge groups vanish in these models¹⁰. Thus, we have $\mathcal{N} = 2$ superconformal gauge theories living in the world-volumes of the D3-branes¹¹.

B. Orbifolds of Odd Order

Let us now discuss the ΩJ orientifold of Type IIB on $\mathcal{M} = \mathbf{C}^3/\Gamma$, where the orbifold group $\Gamma \approx \mathbf{Z}_M$ has odd order ($M \in 2\mathbf{N} + 1$). Such an orientifold contains an arbitrary number of D3-branes. The number of D7-branes, however, is now zero which follows from the untwisted tadpole cancellation conditions.

To understand the twisted tadpole cancellation conditions, first consider $\Omega\tilde{J}$ orientifold of Type IIB on $\mathcal{M}$ (for odd M). This theory has only D7-branes and no D3-branes. The twisted tadpole cancellation conditions for the ΩJ orientifold of Type IIB on $\mathcal{M}$ are isomorphic (after interchanging the corresponding D3- and D7-brane Chan-Paton matrices) to those for the $\Omega\tilde{J}$ orientifold of Type IIB on $\mathcal{M}$ provided that the tadpole cancellation conditions for the $\Omega\tilde{J}$ orientifold of Type IIB on $\mathcal{M}' = \mathbf{C}^3/\Gamma'$ can be satisfied, where $\Gamma' = \{g^k, Rg^k | k = 0, \dots, M-1\} \approx \mathbf{Z}_{2M}$. If the latter condition is not satisfied, then there is no solution to the twisted tadpole cancellation conditions for the ΩJ orientifold of Type IIB on $\mathcal{M}$. The discussion in the previous subsection then implies that the twisted tadpole cancellation conditions can be satisfied only for the ΩJ orientifold of Type IIB on $\mathcal{M}$ with $M = 3$.

The twisted tadpole cancellation conditions for the $\Omega\tilde{J}$ orientifold of Type IIB on $\mathcal{M}$ (for arbitrary odd M) can be computed following [22], and are given by:

$$\text{Tr}(\gamma_{2k,7}) - 4 \sin^2 \left(\frac{2\pi k}{M} \right) \text{Tr}(\gamma_{2k,3}) - 8 \cos^2 \left(\frac{\pi k}{M} \right) = 0 . \quad (24)$$

(Note that the difference between these tadpole cancellation conditions and those of [22] is that we have $8 \cos^2(\pi k/M)$ instead of $32 \cos^2(\pi k/M)$ for the reasons discussed in the previous subsection.) The above discussion then implies the following solution to the twisted tadpole cancellation conditions for the ΩJ orientifold of Type IIB on $\mathcal{M}$ with $M = 3$ ($N = (n_3 + 2\eta)/3$):

$$\gamma_{1,3} = \text{diag}(\exp(2\pi i/3) \text{ (} N \text{ times)}, \exp(-2\pi i/3) \text{ (} N \text{ times)}, 1 \text{ (} N - 2\eta \text{ times)}) . \quad (25)$$

Here $\eta = -1$ if the Ω projection is of the SO type, and $\eta = +1$ if it is of the Sp type¹². The massless spectra (for both choices of η) of these models are given in Table I.

¹⁰The Abelian gauge couplings run in these models so that $U(1)$'s decouple at low energies.

¹¹The 77 gauge groups for large enough values of N are infrared free. They can therefore be treated as global symmetries (in the context of four dimensional gauge theories living on the D3-branes) at low energies.

¹²In this case we have D3-branes only and (unlike in the cases with even M) there is a choice for the action of Ω on the Chan-paton charges.

Note that the one-loop β -functions for the non-Abelian subgroups of the 33 open string sector gauge groups vanish in these models¹³. Thus, we have $\mathcal{N} = 2$ superconformal gauge theories living in the world-volumes of the D3-branes.

C. Comments

In this subsection we would like to comment on some of the issues relevant for the $\mathcal{N} = 2$ superconformal models constructed in the previous subsections.

First, note that the world-sheet parity reversal Ω maps the g^k twisted closed string sector to the g^{M-k} twisted closed string sector. This implies that (for $k \neq 0, M/2$) the g^k and g^{M-k} twisted closed string sectors together give rise to one hypermultiplet and one vector multiplet. On the other hand, the $\mathbf{Z}_2$ twisted sector (present for $M \in 2\mathbf{N}$) gives rise to one hypermultiplet only.

The second comment concerns finiteness of $\mathbf{Z}_6$ and $\mathbf{Z}_3$ theories. From our discussion in section III we (at least naively) expect that models where twisted Chan-Paton matrices are not traceless should not be conformal. Note that in the $\mathbf{Z}_2$ and $\mathbf{Z}_4$ models discussed above the twisted Chan-Paton matrices are traceless so it is not surprising that they are finite. On the other hand, some of the twisted Chan-Paton matrices in the $\mathbf{Z}_6$ and $\mathbf{Z}_3$ models are *not* traceless. Thus, it might appear surprising that they are still finite. Apparently, some subtle cancellation has taken place in these models¹⁴.

We can understand the origin of this “accidental” cancellation in the light of recent results obtained in [23]. In [23] it was shown that the six dimensional orientifold of Type IIB on $T^4/\mathbf{Z}_6$ is on the same moduli as the orientifold of Type IIB on $T^4/\mathbf{Z}_2$ with an (untwisted) NS-NS antisymmetric tensor background. The twisted sector Chan-Paton matrices in the $\mathbf{Z}_2$ model are traceless (for turning on the untwisted NS-NS B -field does not affect the twisted tadpole cancellation conditions [23]). Since the four dimensional orientifolds we are considering here are related to the six dimensional ones (upon compactifying on T^2 , T-dualizing and taking the dimensions of the dual $\tilde{T}^2$ to infinity), it is not surprising that the $\mathbf{Z}_6$ model is finite (that is, this explains the “accidental” cancellation mentioned above). The finiteness of the $\mathbf{Z}_3$ model then follows as $\mathbf{Z}_6 \approx \mathbf{Z}_3 \otimes \mathbf{Z}_2$, and the $\mathbf{Z}_2$ model is finite due to tracelessness of the twisted Chan-Paton matrices.

Here we should mention that there are other $\mathcal{N} = 2$ models which are not superconformal. One way of constructing such models is to consider the $\Omega\tilde{J}$ orientifold of Type IIB on $\mathcal{M} = \mathbf{C}^3/\Gamma$, where $\Gamma \approx Z_M$ has odd order ($M \in 2\mathbf{N} + 1$). Note that in these models there is an orientifold 7-plane (which requires presence of 8 D7-branes) but no orientifold 3-plane¹⁵. Nonetheless, we can introduce an arbitrary number of D3-branes. The gauge group

¹³Just as in the previous cases, the Abelian gauge couplings run in these models so that $U(1)$ ’s decouple at low energies.

¹⁴This cancellation need occur at one-loop level only since these models have $\mathcal{N} = 2$ supersymmetry.

¹⁵These models are “T-dual” (in the non-compact limit) to some of the six dimensional models

coming from the D3-branes is (generically) a product of unitary subgroups, and contains bi-fundamental matter just as in [5,1]. However, there is additional matter coming from the 37 open string sector (which gives fundamentals in the 33 gauge group which also transform as fundamentals under the 77 gauge group). The twisted tadpole cancellation conditions are given by (24). It is not difficult to show that in these models (the non-Abelian part of) the 33 gauge theory is not superconformal (and, in particular, the corresponding one-loop β -functions are non-vanishing). Note that this is in accord with expectations in section III. (The “accidental” finiteness of the gauge theories discussed in the previous subsections is therefore a special feature of those models.) It would be interesting to study these non-finite gauge theories further in the present string theory framework for they may bring additional insight relevant for understanding more general non-conformal gauge theories.

V. $\mathcal{N} = 1$ GAUGE THEORIES

In this section we construct a four dimensional $\mathcal{N} = 1$ gauge theory which has vanishing β -function to all orders in perturbation theory in the large N limit. This particular gauge theory is obtained by appropriately orientifolding Type IIB on $\mathbf{C}^3/(\mathbf{Z}_2 \otimes \mathbf{Z}_2)$. We will also discuss orientifolds corresponding to other orbifold groups (which are subgroups of $SU(3)$) that lead to $\mathcal{N} = 1$ gauge theories which are *not* superconformal.

A. The $\mathbf{Z}_2 \otimes \mathbf{Z}_2$ Model

We start from Type IIB string theory on $\mathcal{M} = \mathbf{C}^3/\Gamma$, where $\Gamma = \{1, R_1, R_2, R_3\} \approx \mathbf{Z}_2 \otimes \mathbf{Z}_2$ ($R_i R_j = R_k$, $i \neq j \neq k \neq i$) is the orbifold group whose action on the complex coordinates z_i ($i = 1, 2, 3$) on $\mathbf{C}^3$ is given by $R_i z_j = -(-1)^{\delta_{ij}} z_j$. Next, we consider an orientifold of this theory where the orientifold action is given by ΩJ . The orientifold group is given by $\mathcal{O} = \{1, R_1, R_2, R_3, \Omega J, \Omega R_1, \Omega R_2, \Omega R_3\}$.

This model is a “T-dual” (in the non-compact limit) of the model studied in [25]. The untwisted tadpole cancellation conditions require presence of three sets of D7-branes with 8 D7-branes in each set. Thus, the locations of D7_{*i*}-branes are given by points in the z_i complex plane. The number of D3-branes is unconstrained (which is due to the fact that the space transverse to the D3-branes is non-compact). The twisted tadpole cancellation conditions imply that the corresponding Chan-Paton matrices $\gamma_{R_i,3}$ and $\gamma_{R_i,7_j}$ are traceless:

$$\text{Tr}(\gamma_{R_i,3}) = \text{Tr}(\gamma_{R_i,7_j}) = 0 . \quad (26)$$

A choice¹⁶ consistent with requirements that the Chan-Paton matrices form a (projective) representation of (the double cover) of Γ is given by ($N = n_3/2$)

$$\gamma_{R_i,3} = i\sigma_i \otimes \mathbf{I}_N , \quad (27)$$

discussed in [24].

¹⁶This choice is unique up to equivalent representations [25].

where σ_i are Pauli matrices, and $\mathbf{I}_N$ is an $N \times N$ identity matrix. (The action on the $D7_i$ Chan-Paton charges is similar.) The spectrum of this model is given in Table II.

Note that the one-loop β -function for the 33 open string sector gauge group vanishes in this model¹⁷. Moreover, all the twisted Chan-Paton matrices are traceless in this model. Following our discussion in section III we thus conclude that this four dimensional gauge theory is superconformal (to all loop orders) in the large N limit.

For completeness, let us give the tree-level superpotential for this model. Let Φ_i , Φ_j^i , Q^i and Q^{ij} be the matter fields in the 33, $7_i 7_i$, 37_i and $7_i 7_j$ open string sectors, respectively. The subscript in Φ_i and Φ_j^i labels three different chiral superfields (see Table II) in the 33 and $7_i 7_i$ sectors. The superpotential can be computed as in [25] and is given by (here we suppress the actual values of the Yukawa couplings)

$$\mathcal{W} = \epsilon_{ijk} \Phi_i \Phi_j \Phi_k + \epsilon_{ijk} \Phi_i^l \Phi_j^l \Phi_k^l + \epsilon_{ijk} \Phi_k^i Q^{ij} Q^{ij} + \Phi_i Q^i Q^i + Q^{ij} Q^{jk} Q^{ki} + Q^{ij} Q^i Q^j . \quad (28)$$

Here the summation over repeated indices is understood.

B. Other $\mathcal{N} = 1$ Gauge Theories from Orientifolds

In this subsection we consider $\mathcal{N} = 1$ orientifolds with orbifold groups such that twisted Chan-Paton matrices are *not* traceless. Unlike the $\mathcal{N} = 2$ examples (where certain “accidental” cancellations at one-loop order led to finiteness as discussed in the previous section) the $\mathcal{N} = 1$ models no longer have vanishing one-loop β -functions (as expected from our discussion in section III).

Instead of being most general, for illustrative purposes in this subsection we will confine our attention to ΩJ orientifolds of Type IIB on $\mathbf{C}^3/\Gamma$ where the orbifold group $\Gamma = \{g^k | k = 0, \dots, M-1\} \approx \mathbf{Z}_M$ (M is odd) is a subgroup of $SU(3)$ (but not of $SU(2)$). The action of Γ on the complex coordinates z_i ($i = 1, 2, 3$) on $\mathbf{C}^3$ is given by $gz_i = \omega^{\ell_i} z_i$, where $\omega = \exp(2\pi i/M)$, $\ell_i \neq 0$, and $\ell_1 + \ell_2 + \ell_3 = M$.

In these orientifolds we have D3-branes only whose number is arbitrary. The twisted tadpole cancellation conditions for these orientifolds are isomorphic (upon interchanging the corresponding D3- and D9-brane Chan-Paton matrices) to those for the Ω orientifolds of Type IIB on $\mathbf{C}^3/\Gamma$. The latter tadpole cancellation conditions were derived in [26]. Applying those results to the cases under consideration we have the following twisted tadpole cancellation conditions ($k = 1, \dots, N-1$):

$$\text{Tr}(\gamma_{2k,3}) = -4\eta \prod_{i=1}^3 (1 + \omega^{k\ell_i}) . \quad (29)$$

Here $\eta = -1$ if the Ω projection is of the SO type, and $\eta = +1$ if it is of the Sp type.

It is not difficult to solve these tadpole cancellation conditions for the general case. For illustrative purposes, however, here we will discuss only two particular examples.

¹⁷For $N > 4$ the $7_i 7_i$ gauge groups are infrared free. They can therefore be treated as global symmetries (in the context of the four dimensional gauge theory living on the D3-branes) at low energies.

First consider the case where $M = 3$. This model is a “T-dual” (in the non-compact limit) of the model studied in [27]¹⁸. The solution to the twisted tadpole cancellation conditions reads ($N = (n_3 - 4\eta)/3$):

$$\gamma_{1,3} = \text{diag}(\exp(2\pi i/3) \text{ (} N \text{ times)}, \exp(-2\pi i/3) \text{ (} N \text{ times)}, 1 \text{ (} N + 4\eta \text{ times)}) . \quad (30)$$

The massless spectra (for both choices of η) of these models are given in Table III. The non-Abelian gauge anomaly is cancelled in this model. However, there is an anomalous $U(1)$. (The superpotential for this model can be found in [27,30].)

Next, let us consider the case where $M = 7$. This model is a “T-dual” (in the non-compact limit) of the model studied in [26]. The solution to the twisted tadpole cancellation conditions reads ($N = (n_3 + 4\eta)/7$):

$$\begin{aligned} \gamma_{1,3} = \text{diag} & (\exp(2\pi i/7) \text{ (} N \text{ times)}, \exp(-2\pi i/7) \text{ (} N \text{ times)}, \\ & \exp(4\pi i/7) \text{ (} N \text{ times)}, \exp(-4\pi i/7) \text{ (} N \text{ times)}, \\ & \exp(6\pi i/7) \text{ (} N \text{ times)}, \exp(-6\pi i/7) \text{ (} N \text{ times)}, 1 \text{ (} N - 4\eta \text{ times)}) . \end{aligned} \quad (31)$$

The massless spectra (for both choices of η) of these models are given in Table III. The non-Abelian gauge anomaly is cancelled in this model. However, there is an anomalous $U(1)$. (The superpotential for this model can be found in [26].)

Note that none of the above two models have vanishing one-loop β -functions for the non-Abelian factors in the gauge group. Moreover, the presence of anomalous $U(1)$ [31] implies that there is a Fayet-Iliopoulos D-term which must be cancelled via a generalized Green-Schwarz mechanism [32]. Thus, some fields must acquire non-zero vevs to cancel the Fayet-Iliopoulos D-term. Using Type I-Heterotic duality [29] it was argued in [30] that the twisted closed string sector states transform non-trivially under the anomalous $U(1)$ gauge transformation, and (together with the dilaton plus axion supermultiplet) can cancel the Fayet-Iliopoulos D-term. In this process a non-zero vev is generated for the field responsible for this cancellation.

Thus, in the $\mathcal{N} = 1$ cases we see that the models with non-vanishing traces of the twisted Chan-Paton matrices lead (as expected in section III) to non-finite models. In particular, the “accidental” vanishing of the one-loop β -functions (as in the corresponding $\mathcal{N} = 2$ cases) does not occur.

Here we should mention that the tadpole cancellation condition (29) is necessary not only for ultraviolet finiteness of the corresponding theories but also for non-Abelian gauge anomaly cancellation. For illustrative purposes, to see what can go wrong if we relax this condition, let us consider the above $\mathbf{Z}_3$ example with the following choice for the twisted Chan-Paton matrices:

$$\gamma_{1,3} = \text{diag}(\exp(2\pi i/3) \text{ (} N \text{ times)}, \exp(-2\pi i/3) \text{ (} N \text{ times)}, 1 \text{ (} N' \text{ times)}) . \quad (32)$$

The gauge group in the 33 open string sector is $U(N) \otimes G_\eta(N')$, and the chiral matter is given by $3(\mathbf{R}_\eta, \mathbf{1})$ and $3(\overline{\mathbf{N}}, \mathbf{N}')$ (see Table III for notation). Note that $\mathbf{R}_\eta$ of $SU(N)$

¹⁸Also see, *e.g.*, [28].

contributes as much as $N + 4\eta$ chiral superfields in $\mathbf{N}$ of $SU(N)$ into the non-Abelian gauge anomaly. Thus, the non-Abelian gauge anomaly cancellation implies that $N' = N + 4\eta$. This is precisely the solution to the tadpole cancellation condition (29).

C. Cases without World-Sheet Description

In subsection C of section III we mentioned that for certain choices of the orbifold group the corresponding orientifolds may not have a world-sheet description. Here we give an explicit example of this.

Consider Type IIB on $\mathbf{C}^3/\Gamma$ with $\Gamma \approx D_N$ (non-Abelian dihedral group) where the action of Γ on the complex coordinates $z_i (i = 1, 2, 3)$ on $\mathbf{C}^3$ is given by ($\omega = \exp(2\pi i/N)$):

$$gz_1 = z_1, \quad gz_2 = \omega z_2, \quad gz_3 = \omega^{-1} z_3, \quad (33)$$

$$rz_1 = -z_1, \quad rz_2 = z_3, \quad rz_3 = z_2, \quad (34)$$

where g, r are the generators of D_N . Note that g and r do not commute: $rg = g^{-1}r$.

Now consider ΩJ ($Jz_i = -z_i$) orientifold of this theory. The orientifold group is $\mathcal{O} = \{g^k, rg^k, \Omega Jg^k, \Omega Jr g^k | k = 0, \dots, N-1\}$. Note that $(Jrg^k)^2 = 1$, and the set of points in $\mathbf{C}^3$ fixed under the action of Jrg^k has real dimension two. This implies that there are N kinds of orientifold 7-planes corresponding to the elements $\Omega Jr g^k$. Note, however, that due to non-commutativity between g and r (and, therefore, between different Jrg^k), these orientifold 7-planes (as well as the corresponding D7-branes) are mutually non-local. This implies that this orientifold does not have a world-sheet description. One way to understand such models (at least in compact cases) is to use their F-theory [33] description.

Here we should mention that there are other models with both Abelian and non-Abelian orbifold groups for which the perturbative description in terms of open strings stretched between mutually local D-branes is inadequate. We refer the reader to [18] for details.

ACKNOWLEDGMENTS

I would like to thank Michael Bershadsky, Andrei Johansen, Henry Tye, and especially Cumrun Vafa for valuable discussions. This work is supported in part by the grant NSF PHY-96-02074, and the DOE 1994 OJI award. I would also like to thank Albert and Ribena Yu for financial support.

FIGURES





FIG. 1. A planar diagram.



Klein bottle




Möbius strip




Annulus


FIG. 2. Tree-channel Klein bottle, Möbius strip and annulus amplitudes.



Two cross-caps



A cross-cap and a D-brane



Two D-branes

FIG. 3. Factorization of the Klein bottle, Möbius strip and annulus amplitudes.

TABLES

Model	Gauge Group	Charged Hypermultiplets	Twisted Sector Hypermultiplets	Twisted Sector Vector Multiplets
$\mathbf{Z}_2$	$U(N)_{33} \otimes U(4)_{77}$	$2(\mathbf{A}; \mathbf{1})_{33}$ $2(\mathbf{1}; \mathbf{6})_{77}$ $(\mathbf{N}; \mathbf{4})_{37}$	1	0
$\mathbf{Z}_4$	$[U(N) \otimes U(N)]_{33} \otimes$ $[U(2) \otimes U(2)]_{77}$	$(\mathbf{A}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{33}$ $(\mathbf{1}, \mathbf{A}; \mathbf{1}, \mathbf{1})_{33}$ $(\mathbf{N}, \mathbf{N}; \mathbf{1}, \mathbf{1})_{33}$ $(\mathbf{1}, \mathbf{1}; \mathbf{1}', \mathbf{1})_{77}$ $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1}')_{77}$ $(\mathbf{1}, \mathbf{1}; \mathbf{2}, \mathbf{2})_{77}$ $(\mathbf{N}, \mathbf{1}; \mathbf{2}, \mathbf{1})_{37}$ $(\mathbf{1}, \mathbf{N}; \mathbf{1}, \mathbf{2})_{37}$	2	1
$\mathbf{Z}_6$	$[U(N) \otimes U(N) \otimes U(N+1)]_{33} \otimes$ $[U(1) \otimes U(1) \otimes U(2)]_{77}$	$(\mathbf{A}, \mathbf{1}, \mathbf{1}; \mathbf{1})_{33}$ $(\mathbf{1}, \mathbf{A}, \mathbf{1}; \mathbf{1})_{33}$ $(\mathbf{N}, \mathbf{1}, \mathbf{N} + \mathbf{1}; \mathbf{1})_{33}$ $(\mathbf{1}, \mathbf{N}, \mathbf{N} + \mathbf{1}; \mathbf{1})_{33}$ $(\mathbf{1}, \mathbf{1}, \mathbf{1}; \mathbf{2}_1)_{77}$ $(\mathbf{1}, \mathbf{1}, \mathbf{1}; \mathbf{2}_2)_{77}$ $(\mathbf{N}, \mathbf{1}, \mathbf{1}; \mathbf{1}_1)_{37}$ $(\mathbf{1}, \mathbf{N}, \mathbf{1}; \mathbf{1}_2)_{37}$ $(\mathbf{1}, \mathbf{1}, \mathbf{N} + \mathbf{1}; \mathbf{2})_{37}$	3	2
$\mathbf{Z}_3$	$[U(N) \otimes G_\eta(N - 2\eta)]_{33}$	$(\mathbf{R}_\eta, \mathbf{1})_{33}$ $(\mathbf{N}, \mathbf{N} - \mathbf{2}\eta)_{33}$	1	1

TABLE I. The massless spectra of $\mathcal{N} = 2$ orientifolds of Type IIB on $\mathbf{C}^3/\mathbf{Z}_N$ $M = 2, 3, 4, 6$. The semi-colon in the column “Charged Hypermultiplets” separates 33 and 77 representations. The notation $\mathbf{A}$ stands for the two-index antisymmetric representation (which is $N(N-1)/2$ dimensional for $U(N)$) of the corresponding unitary group. The notation $\mathbf{1}'$ in the $\mathbf{Z}_4$ model stands for the antisymmetric representation of the corresponding $U(2)$ group (whose charge with respect to the $U(1)$ factor is 2). In the $\mathbf{Z}_6$ model the subscripts “1” and “2” indicate that the corresponding states are charged under the first and the second $U(1)$ factors in the 77 gauge group. In the $\mathbf{Z}_3$ model $G_\eta = SO$ for $\eta = -1$ and $G_\eta = Sp$ for $\eta = +1$. (Here we are using the convention that $Sp(2m)$ has rank m .) Also, $R_\eta = \mathbf{A}$ for $\eta = -1$, and $R_\eta = \mathbf{S}$ (two-index $N(N+1)/2$ dimensional symmetric representation of $U(N)$) for $\eta = +1$. By twisted sector hypermultiplets/vector multiplets we mean those in the twisted *closed* string sectors. The untwisted closed string sector states are not shown.

Model	Gauge Group	Charged Chiral Multiplets	Twisted Sector Chiral Multiplets	Twisted Sector Vector Multiplets
$\mathbf{Z}_2 \otimes \mathbf{Z}_2$	$Sp(N)_{33} \otimes$ $Sp(4)_{7_1 7_1} \otimes Sp(4)_{7_2 7_2} \otimes Sp(4)_{7_3 7_3}$	$3(\mathbf{A}; \mathbf{1}; \mathbf{1}; \mathbf{1})_{33}$ $3(\mathbf{1}; \mathbf{6}; \mathbf{1}; \mathbf{1})_{7_1 7_1}$ $3(\mathbf{1}; \mathbf{1}; \mathbf{6}; \mathbf{1})_{7_2 7_2}$ $3(\mathbf{1}; \mathbf{1}; \mathbf{1}; \mathbf{6})_{7_3 7_3}$ $(\mathbf{N}; \mathbf{4}; \mathbf{1}; \mathbf{1})_{37_1}$ $(\mathbf{N}; \mathbf{1}; \mathbf{4}; \mathbf{1})_{37_2}$ $(\mathbf{N}; \mathbf{1}; \mathbf{1}; \mathbf{4})_{37_3}$ $(\mathbf{1}; \mathbf{4}; \mathbf{4}; \mathbf{1})_{7_1 7_2}$ $(\mathbf{1}; \mathbf{1}; \mathbf{4}; \mathbf{4})_{7_2 7_3}$ $(\mathbf{1}; \mathbf{4}; \mathbf{1}; \mathbf{4})_{7_3 7_1}$	3	0

TABLE II. The massless spectrum of $\mathcal{N} = 1$ orientifold of Type IIB on $\mathbf{C}^3/(\mathbf{Z}_2 \otimes \mathbf{Z}_2)$. The notation $\mathbf{A}$ stands for the two-index antisymmetric (reducible) representation of $Sp(N)$. The untwisted closed string sector states are not shown.

Model	Gauge Group	Charged Chiral Multiplets	Twisted Sector Chiral Multiplets	Twisted Sector Vector Multiplets
$\mathbf{Z}_3$	$[U(N) \otimes G_\eta(N + 4\eta)]_{33}$	$3(\mathbf{R}_\eta, \mathbf{1})(+2)_{33}$ $3(\overline{\mathbf{N}}, \mathbf{N} + 4\eta)(-1)_{33}$	1	0
$\mathbf{Z}_7$	$[U(N) \otimes U(N) \otimes U(N) \otimes$ $G_\eta(N - 4\eta)]_{33}$	$(\mathbf{R}_\eta, \mathbf{1}, \mathbf{1}, \mathbf{1})(+2, 0, 0)_{33}$ $(\mathbf{1}, \mathbf{R}_\eta, \mathbf{1}, \mathbf{1})(0, +2, 0)_{33}$ $(\mathbf{1}, \mathbf{1}, \mathbf{R}_\eta, \mathbf{1})(0, 0, +2)_{33}$ $(\mathbf{N}, \mathbf{1}, \mathbf{1}, \mathbf{N} - 4\eta)(+1, 0, 0)_{33}$ $(\mathbf{1}, \mathbf{N}, \mathbf{1}, \mathbf{N} - 4\eta)(0, +1, 0)_{33}$ $(\mathbf{1}, \mathbf{1}, \mathbf{N}, \mathbf{N} - 4\eta)(0, 0, +1)_{33}$ $(\overline{\mathbf{N}}, \mathbf{N}, \mathbf{1}, \mathbf{1})(-1, +1, 0)_{33}$ $(\mathbf{1}, \overline{\mathbf{N}}, \mathbf{N}, \mathbf{1})(0, -1, +1)_{33}$ $(\mathbf{N}, \mathbf{1}, \overline{\mathbf{N}}, \mathbf{1})(+1, 0, -1)_{33}$ $(\overline{\mathbf{N}}, \overline{\mathbf{N}}, \mathbf{1}, \mathbf{1})(-1, -1, 0)_{33}$ $(\mathbf{1}, \overline{\mathbf{N}}, \overline{\mathbf{N}}, \mathbf{1})(0, -1, -1)_{33}$ $(\overline{\mathbf{N}}, \mathbf{1}, \overline{\mathbf{N}}, \mathbf{1})(-1, 0, -1)_{33}$	3	0

TABLE III. The massless spectra of $\mathcal{N} = 1$ orientifolds of Type IIB on $\mathbf{C}^3/\mathbf{Z}_M$ ($M = 3, 7$). Here we are using some of the notations from Table I. The $U(1)$ charges of the states in the 33 open string sector are given in parentheses. The untwisted closed string sector states are not shown.

REFERENCES

- [1] M. Bershadsky, Z. Kakushadze and C. Vafa, “String Expansion as Large N Expansion of Gauge Theories”, hep-th/9803076.
- [2] G. ‘t Hooft, “A Planar Diagram Theory For Strong Interactions”, Nucl. Phys. **B72** (1974) 461.
- [3] E. Witten, “Chern-Simons Gauge Theory As A String Theory”, hep-th/9207094.
- [4] S. Kachru and E. Silverstein, “4d Conformal Field Theories and Strings on Orbifolds”, hep-th/9802183.
- [5] A. Lawrence, N. Nekrasov and C. Vafa, “On Conformal Theories in Four Dimensions”, hep-th/9803015.
- [6] L.E. Ibáñez, “A Chiral D=4, N=1 String Vacuum with a Finite Low Energy Effective Field Theory”, hep-th/9802103.
- [7] A. Hanany, M.J. Strassler and A.M. Uranga, “Finite Theories and Marginal Operators on the Brane”, hep-th/9803086.
- [8] D.A. Kosower, B.-H. Lee and V.P. Nair, “Multi Gluon Scattering: A String Based Calculation”, Phys. Lett. **B201** (1988) 85;
Z. Bern and D.A. Kosower, “Efficient Calculation of One Loop QCD Amplitudes”, Phys. Rev. Lett. **66** (1991) 1669.
- [9] Z. Bern, L. Dixon, D.C. Dunbar, M. Perelstein and J.S. Rozowsky, “On the Relationship between Yang-Mills Theory and Gravity and Its Implication for Ultraviolet Divergences”, hep-th/9802162.
- [10] I.R. Klebanov, “World Volume Approach to Absorption by Non-dilatonic Branes”, Nucl. Phys. **B496** (1997) 231, hep-th/9702076.
- [11] S.S. Gubser and I.R. Klebanov, “Absorption by Branes and Schwinger Terms in the World Volume Theory”, Phys. Lett. **B413** (1997) 41, hep-th/9708005.
- [12] J.M. Maldacena, “The Large N Limit of Superconformal Field Theories and Supergravity”, hep-th/9711200.
- [13] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge Theory Correlators from Non-Critical String Theory”, hep-th/9802109.
- [14] G.T. Horowitz and H. Ooguri, “Spectrum of Large N Gauge Theory from Supergravity”, hep-th/9802116.
- [15] E. Witten, “Anti-de Sitter Space And Holography”, hep-th/9802150; “Anti-de Sitter Space, Thermal Phase Transition, And Confinement In Gauge Theories”, hep-th/9803131.
- [16] S.S. Gubser, I.R. Klebanov and A.W. Peet, “Entropy and Temperature of Black 3-branes”, Phys. Rev. **D54** (1996) 3915, hep-th/9602135;
M.R. Douglas, J. Polchinski and A. Strominger, “Probing Five Dimensional Black Holes with D-branes”, hep-th/9703031;
S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, “String Theory and Classical Absorption by Three Branes”, Nucl. Phys. **B499** (1997) 217, hep-th/9703040;
J.M. Maldacena and A. Strominger, “Semiclassical Decay of Near Extremal Fivebranes”, hep-th/9710014;
A.M. Polyakov, “String Theory and Quark Confinement”, hep-th/9711002;
K. Sfetsos and K. Skenderis, “Microscopic Derivation of the Bekenstein-Hawking Entropy Formula for Non-extremal Black Holes”, hep-th/9711138;

S. Ferrara and C. Fronsdal, “Conformal Maxwell Theory as a Singleton Field Theory on AdS_5 , IIB Three Branes and Duality”, hep-th/9712239;

N. Itzhaki, J.M. Maldacena, J. Sonnenschein and S. Yankielowicz, “Supergravity and the Large N Limit of Theories With Sixteen Supercharges”, hep-th/9802042;

M. Gunaydin and D. Minic, “Singletons, Doubletons and M Theory”, hep-th/9802047;

M. Berkooz, “A Supergravity Dual of a (1,0) Field Theory in Six Dimensions”, hep-th/9802195;

V. Balasubramanian and F. Larsen, “Near Horizon Geometry and Black Holes in Four Dimensions”, hep-th/9802198;

S. Ferrara, C. Fronsdal and A. Zaffaroni, “On $N=8$ Supergravity on AdS_5 and $N=4$ Superconformal Yang-Mills theory”, hep-th/9802203;

S.-J. Rey and J.-T. Yee, “Macroscopic Strings as Heavy Quarks of Large N Gauge Theory and Anti-de Sitter Supergravity”, hep-th/9803001;

J.M. Maldacena, “Wilson loops in large N field theories”, hep-th/9803002;

M. Flato and C. Fronsdal, “Intersecting Singletons”, hep-th/9803013;

S.S. Gubser, A. Hashimoto, I.R. Klebanov and M. Krasnitz, “Scalar Absorption and the Breaking of the Worldvolume Conformal Invariance”, hep-th/9803023;

I.Ya. Aref’eva and I.V. Volovich, “On Large N Conformal Theories, Field Theories in the Anti-de Sitter space and Singletons”, hep-th/9803023;

L. Castellani, A. Ceresole, R. D’Auria, S. Ferrara, P. Fré and M. Trigiante, “ G/H M-branes and AdS_{p+2} Geometries”, hep-th/9803039;

O. Aharony, Y. Oz and Z. Yin, “M Theory on $AdS_p \times S^{11-p}$ and Superconformal Field Theories”, hep-th/9803051;

S. Minwala, “Particles on $AdS_{4/7}$ and Primary Operators on $M_{2/5}$ Brane Worldvolumes”, hep-th/9803053;

S. Ferrara and A. Zaffaroni, “ $N=1,2$ 4D Superconformal Field Theories and Supergravity in AdS_5 ”, hep-th/9803060;

M.J. Duff, H. Lu and C.N. Pope, “ $AdS_5 \times S^5$ Untwisted”, hep-th/9803061;

R.G. Leigh and M. Rozali, “The Large N Limit of the (2,0) Superconformal Field Theory”, hep-th/9803068;

E. Halyo, “Supergravity on $AdS_{4/7} \times S^{7,4}$ and M Branes”, hep-th/9803077; “Supergravity on $AdS_{5/4} \times$ Hopf Fibrations and Conformal Field Theories”, hep-th/9803193;

A. Rajaraman, “Two-Form Fields and the Gauge Theory Description of Black Holes”, hep-th/9803082;

G.T. Horowitz and S.F. Ross, “Possible Resolution of Black Hole Singularities from Large N Gauge Theory”, hep-th/9803085;

S. Ferrara, A. Kehagias, H. Partouche and A. Zaffaroni, “Membranes and Fivebranes with Lower Supersymmetry and their AdS Supergravity Duals”, hep-th/9803109;

J.A. Minahan, “Quark-Monopole Potentials in Large N Super Yang-Mills”, hep-th/9803111;

J. Gomis, “Anti de Sitter Geometry and Strongly Coupled Gauge Theories”, hep-th/9803019;

S.-J. Rey, S. Theisen and J.-T. Yee, “Wilson-Polyakov Loop at Finite Temperature in Large N Gauge Theory and Anti-de Sitter Supergravity”, hep-th/9803135;

A. Brandhuber, N. Itzhaki, J. Sinnenschein and S. Yankielowicz, “Wilson Loops in the

- Large N Limit at Finite Temperature”, hep-th/9803137;
M. Gunaydin, “Unitary Supermultiplets of $\text{OSp}(1/32, \text{R})$ and M-theory”, hep-th/9803138;
Y. Oz and J. Terning, “Orbifolds of $AdS_5 \times S^5$ and 4d Conformal Field Theories”, hep-th/9803167;
L. Andrianopoli and S. Ferrara, “K-K excitations on $AdS_5 \times S^5$ as N=4 ‘primary’ superfields”, hep-th/9803171;
I.V. Volovich, “Large N Gauge Theories and Anti-de Sitter Bag Model”, hep-th/9803174;
F. Brandt, J. Gomis and J. Simon, “D-string on Near Horizon Geometries and Infinite Conformal Symmetry”, hep-th/9803196.
- [17] V. Kaplunovsky, “One-Loop Threshold Effects in String Unification”, Nucl. Phys. **B307** (1988) 145; ERRATA for “One-Loop Threshold Effects in String Unification”, Nucl. Phys. **B382** (1992) 436, hep-th/9205068.
 - [18] Z. Kakushadze, G. Shiu and S.-H.H. Tye, “Type IIB Orientifolds, F-theory, Type I Strings on Orbifolds and Type I - Heterotic Duality”, preprint CLNS 98/1549, HUTP-97/A044, NUB 3168, to appear.
 - [19] G. Zwart, “Four-dimensional $N = 1$ $Z_N \times Z_M$ Orientifolds”, hep-th/9708040.
 - [20] G. Pradisi and A. Sagnotti, “Open String Orbifolds”, Phys. Lett. **B216** (1989) 59;
M. Bianchi and A. Sagnotti, “On the Systematics of Open String Theories”, Phys. Lett. **B247** (1990) 517; “Twist Symmetry and Open String Wilson Lines”, Nucl. Phys. **B361** (1991) 519.
 - [21] E.G. Gimon and J. Polchinski, “Consistency Conditions for Orientifolds and D-Manifolds”, Phys. Rev. **D54** (1996) 1667, hep-th/9601038.
 - [22] E.G. Gimon and C.V. Johnson, “K3 Orientifolds” Nucl. Phys. **B477** (1996) 715, hep-th/9604129;
A. Dabholkar and J. Park, “Strings on Orientifolds”, Nucl. Phys. **B477** (1996) 701, hep-th/9604178.
 - [23] Z. Kakushadze, G. Shiu and S.-H.H. Tye, “Type IIB Orientifolds with NS-NS Antisymmetric Tensor Backgrounds”, hep-th/9803141.
 - [24] J.D. Blum and K. Intriligator, “Consistency Conditions for Branes at Orbifold Singularities”, Nucl. Phys. **B506** (1997) 223, hep-th/9705030; “New Phases of String Theory and 6d RG Fixed Points via Branes at Orbifold Singularities”, Nucl. Phys. **B506** (1997) 199, hep-th/9705044;
K. Intriligator, “New String Theories in Six Dimensions via Branes at Orbifold Singularities”, hep-th/9708117.
 - [25] M. Berkooz and R.G. Leigh, “A D=4 N=1 Orbifold of Type I Strings”, Nucl. Phys. **B483** (1997) 187, hep-th/9605049.
 - [26] Z. Kakushadze and G. Shiu, “A Chiral $N = 1$ Type I Vacuum in Four Dimensions and Its Heterotic Dual”, Phys. Rev. **D56** (1997) 3686, hep-th/9705163; “4D Chiral $N = 1$ Chiral Vacua with and without $D5$ -branes”, hep-th/9706051.
 - [27] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Ya.S. Stanev, “Chiral Asymmetry in Four-Dimensional Open-String Vacua”, Phys. Lett. **B385** (1996) 96, hep-th/9606169.
 - [28] J. Lykken, E. Poppitz and S. Trivedi, “M(ore) on Chiral Gauge Theories from D-branes”,

- hep-th/9712193.
- [29] J. Polchinski and E. Witten, “Evidence for Heterotic - Type I String Duality”, Nucl. Phys. **B460** (1996) 525, hep-th/9510169.
 - [30] Z. Kakushadze, “Aspects of $N = 1$ Type I-Heterotic Duality in Four Dimensions”, Nucl. Phys. **B512** (1998) 221, hep-th/9704059.
 - [31] E. Witten, “Some Properties of $O(32)$ Superstrings”, Phys. Lett. **B149** (1984) 351;
M. Dine, N. Seiberg and E. Witten, “Fayet-Iliopoulos Terms in String Theory”, Nucl. Phys. **B289** (1987) 589.
 - [32] M. Green and J.H. Schwarz, “Anomaly Cancellations In Supersymmetric D=10 Gauge Theory Require $SO(32)$ ”, Phys. Lett. **B149** (1984) 117; “Infinity Cancellations in $SO(32)$ Superstring Theory”, Phys. Lett. **B151** (1985) 21.
 - [33] C. Vafa, “Evidence for F-theory”, Nucl. Phys. **B469** (1996) 403, hep-th/9602022.